


=> d que
L39



VAR G1=16/18
NODE ATTRIBUTES:
CONNECT IS E1 RC AT 9
CONNECT IS E1 RC AT 16
CONNECT IS E1 RC AT 19
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 15

STEREO ATTRIBUTES: NONE
L41 14 SEA FILE=REGISTRY SSS FUL L39
L42 5 SEA FILE=REGISTRY ABB=ON PLU=ON L41 AND "15"
L43 10 SEA FILE=CAPLUS ABB=ON PLU=ON L42

=> d bib ab hitstr 1-10

L43 ANSWER 1 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2005:426567 CAPLUS
DN 142:482029
TI Preparation of [(1R,3S)-3-isopropyl-3-[[3-(trifluoromethyl)-7,8-dihydro-1,6-naphthyridin-6(5H)-yl]carbonyl]cyclopentyl][(3S,4S)-3-methoxytetrahydro-2H-pyran-4-yl]amine salt as chemokine receptor CCR-2 antagonist
IN Cai, Dongwei; Fleitz, Fred; Ge, Min; Hoerrner, Scott; Javadi, Gary; Jensen, Mark; Larsen, Robert; Li, Wenjie; Nelson, Dorian; Szumigala, Elizabeth; Yang, Lihu; Zhou, Changyou
PA Merck & Co., Inc., USA
SO PCT Int. Appl., 33 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005044795	A1	20050519	WO 2004-US35294	20041025
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

PRAI US 2003-514754P P 20031027
OS CASREACT 142:482029; MARPAT 142:482029
AB The present invention provides an efficient synthesis for the preparation of [(1R,3S)-3-isopropyl-3-[[3-(trifluoromethyl)-7,8-dihydro-1,6-naphthyridin-6(5H)-yl]carbonyl]cyclopentyl][(3S,4S)-3-methoxytetrahydro-2H-pyran-4-yl]amine (I) and its succinate salt. The present invention addnl. provides an efficient syntheses for the preparation of intermediates, i.e. (3R)-3-methoxytetrahydro-4H-pyran-4-one (II), (1S,4S)-4-(2,5-dimethyl-1H-pyrrol-1-yl)-1-isopropylcyclopent-2-ene-1-carboxylic acid (III), and 3-(trifluoromethyl)-5,6,7,8-tetrahydro-1,6-naphthyridine (IV), and for the preparation of the precursor (3S,4S)-N-((1S,4S)-4-isopropyl-4-[[3-(trifluoromethyl)-7,8-dihydro-1,6-naphthyridin-6(5H)-yl]carbonyl]cyclopent-2-en-1-yl)-3-methoxytetrahydro-2H-pyran-4-amine (V). The invention addnl. resides in the superior properties of the I succinate. Thus, 730 g III was treated with 228 mL methanesulfonyl chloride in the presence of 0.93 L diisopropylethylamine in 6.6 L THF at 0-13°, stirred at room temperature for 4 h, cooled to .apprx.15°, treated with IV.HCl, warmed to 23°, treated with 0.93 L diisopropylethylamine with cooling at .apprx.25° over 15 min, and aged for .apprx.1 h to give 1.055 kg the compound (VI). VI (1.055 kg) in MeOH was added to 1 kg hydroxylamine hydrochloride, 50% aqueous hydroxylamine (1 L), and 5 L H2O, and the resulting slurry was refluxed at 71° for 6 h to give, after treatment with anhydrous HCl in isopropanol, the amine dihydrochloride (VII) (0.76 kg). VII (777 g) was slurried in 3 L iso-Pr acetate and the mixture was cooled in an ice bath, successively treated with 860 mL n-Bu3N, 260 mL isopropanol, and sodium triacetoxyborohydride (724 g, at 5°), and after 1 h, treated with a solution of II in iso-Pr acetate (1.76 L of a 160 g/L solution), and allowed to react for 6 h to give 654 g crude V (90%). V (640 g) was diluted with 6.4 L MeOH, charged to an autoclave, treated with a slurry of 256 g

5% Pd-C in 5.0 L MeOH, and hydrogenated under H pressure of 40 psi at 25° overnight to give a solution of 633 g I in MeOH (98.5%) which was concentrated to an oil (770 g). The oil was dissolved in 3.1 L iso-Pr acetate and the solution was concentrated to a brown oil. Dilution with iso-Pr acetate and concentration was repeated two addnl. times. The resulting oil was converted into I benzenesulfonate (1.645 kg) by treating with 283 g benzenesulfonic acid in iso-Pr acetate and treatment with heptane and the resulting benzenesulfonate salt was treated with a mixture of K₂CO₃, H₂O, and iso-Pr acetate to give an oil containing 1.16 kg I. The latter oil was treated with 294 g succinic acid in ethanol and heptane to give 1.328 kg I succinate.

IT 851956-93-9P, Methyl (1S,4S)-4-(acetylamino)-1-isopropylcyclopent-2-ene-1-carboxylate
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of [(1R,3S)-3-isopropyl-3-[[3-(trifluoromethyl)-7,8-dihydro-1,6-naphthyrin-6(5H)-yl]carbonyl]cyclopentyl][(3S,4S)-3-methoxytetrahydro-2H-pyran-4-yl]amine salt as chemokine receptor CCR-2 antagonist)
 RN 851956-93-9 CAPLUS
 CN 2-Cyclopentene-1-carboxylic acid, 4-(acetylamino)-1-(1-methylethyl)-, methyl ester, (1S,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L43 ANSWER 2 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2000:53929 CAPLUS

DN 132:107046

TI Preparation of optically active azabicycloheptenone derivatives by stereospecific enzymic hydrolysis

IN Bernegger-Egli, Christine; Brux, Frank; Roduit, Jean Paul; Werbitzky, Oleg; Guggisberg, Yves

PA Lonza A.-G., Switz.

SO PCT Int. Appl., 27 pp.

CODEN: PIXXD2

DT Patent

LA German

FAN.CNT 1


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000003032	A1	20000120	WO 1999-EP4814	19990708
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9952803	A1	20000201	AU 1999-52803	19990708
EP 1095160	A1	20010502	EP 1999-938217	19990708
EP 1095160	B1	20040128		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
JP 2002520027	T2	20020709	JP 2000-559252	19990708
AT 258605	E	20040215	AT 1999-938217	19990708
PT 1095160	T	20040630	PT 1999-938217	19990708
ES 2214876	T3	20040916	ES 1999-938217	19990708
NO 2001000121	A	20010108	NO 2001-121	20010108
US 6780634	B1	20040824	US 2001-743391	20010417
US 2004167351	A1	20040826	US 2004-779339	20040213
PRAI EP 1998-112719	A	19980709		
EP 1998-123949	A	19981217		
WO 1999-EP4814	W	19990708		
US 2001-743391	A3	20010417		
MARPAT 132:107046				

OS The invention relates to a biotechnol. method for producing optically active compds. of general formulas (I) and (II), wherein R1 represents acyl or acyloxy, and R2 represents H or C1-C10 alkyl, by reaction of the racemic lactam using a hydrolase in the presence of a nucleophile and in the presence of a base in a constant pH range. The invention also relates to the subsequent conversion of compound I into the optically active 1-amino-4-(hydroxymethyl)-2-cyclopentene of formula (III). Racemic 2-acetyl-2-azabicyclo[2.2.1]hept-5-en-3-one 419.25 mL was diluted with water 60 mL and a com. subtilisin solution 31.5 mL. This solution was brought to pH 7.5 and incubated at 30° with vigorous stirring. After 45 h (1R,4S)-2-Acetyl-2-azabicyclo[2.2.1]hept-5-en-3-one with an ee 99% was obtained. Final yield of purified product was 31%.

IT 255839-19-1P
 RL: BMF (Bioindustrial manufacture); BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation of optically active azabicycloheptenone derivs. by stereospecific enzymic hydrolysis)

RN 255839-19-1 CAPLUS
CN 2-Cyclopentene-1-carboxylic acid, 4-(acetlamino)-, propyl ester, (1S,4R)-
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L43 ANSWER 3 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN
AN 1997:480168 CAPLUS
DN 127:176646
TI An approach to a carbocyclic analog of cyclic adenosine 5'-diphosphate ribose. The synthesis and bisphosphorylation of N1-[(1S,3R)-3-(hydroxymethyl)cyclopent-1-yl]inosine
AU Fortt, Simon M.; Potter, Barry V. L.
CS Dep. Medicinal Chem., School Pharmacy and Pharmacology, Univ. Bath, Bath, BA2 7AY, UK
SO Tetrahedron Letters (1997), 38(30), 5371-5374
CODEN: TELEAY; ISSN: 0040-4039
PB Elsevier
DT Journal
LA English
AB The synthesis of N1-[(1S, 3R)-3-(hydroxymethyl)cyclopent-1-yl]inosine via a cyclocondensation route is reported. Regioselective bisphosphorylation of the primary 5'-hydroxyl groups leads to a carbocyclic, inosine analog of seco ADP ribose.
IT 127061-46-5P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and bisphosphorylation of hydroxymethylcyclopentylinosine)
RN 127061-46-5 CAPLUS
CN 2-Cyclopentene-1-carboxylic acid, 4-(acetlamino)-, methyl ester, (1S-cis)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L43 ANSWER 4 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN
AN 1996:569065 CAPLUS
DN 125:300485
TI Lipase-catalyzed resolution of 2-azabicyclo[2.2.1]hept-5-en-3-ones
AU Nakano, Hiroto; Iwasa, Kazuto; Okuyama, Yuko; Hongo, Hiroshi
CS Tohoku College Pharmacy, Sendai, 981, Japan
SO Tetrahedron: Asymmetry (1996), 7(8), 2381-2386
CODEN: TASYE3; ISSN: 0957-4166
PB Elsevier
DT Journal
LA English
AB The lipase-catalyzed asym. resolution of 2-azabicyclo[2.2.1]hept-5-en-3-ones was reported. Non-racemic chiral 2-azabicyclo[2.2.1]hept-5-en-3-ones were obtained conveniently by lipase-catalyzed enantioselective transesterification or hydrolysis of 2-(hydroxymethyl)-2-azabicyclo[2.2.1]hept-5-en-3-one or 2-[(acetyloxy)methyl]-2-azabicyclo[2.2.1]hept-5-en-3-one. The resolution of (+)-2-(hydroxymethyl)-2-azabicyclo[2.2.1]hept-5-en-3-one (I) gave (1R)-2-[(acetyloxy)methyl]-2-azabicyclo[2.2.1]hept-5-en-3-one which was hydrolyzed to give (1R)-2-(hydroxymethyl)-2-azabicyclo[2.2.1]hept-5-en-3-one. Ring opening of the latter gave (-)-(1S-cis)-4-(acetlamino)-2-cyclopentene-1-carboxylic acid Me ester (II) which is an intermediate for carbovir.
IT 127061-46-5P, (1S-cis)-4-(Acetlamino)-2-cyclopentene-1-carboxylic acid methyl ester
RL: SPN (Synthetic preparation); PREP (Preparation)
(lipase-catalyzed resolution of 2-azabicyclo[2.2.1]hept-5-en-3-ones)
RN 127061-46-5 CAPLUS
CN 2-Cyclopentene-1-carboxylic acid, 4-(acetlamino)-, methyl ester, (1S-cis)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L43 ANSWER 5 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1996:175342 CAPLUS
 DN 124:254209
 TI Substrate specificity of glycine ribonucleotide transformylase from chicken liver
 AU Antle, Vincent D.; Liu, Dashan; McKellar, B. Robert; Caperelli, Carol A.; Hua, Mei; Vince, Robert
 CS Division Pharmaceutical Sciences, University Cincinnati Medical Center, Cincinnati, OH, 45267-0004, USA
 SO Journal of Biological Chemistry (1996), 271(11), 6045-9
 CODEN: JBCHA3; ISSN: 0021-9258
 PB American Society for Biochemistry and Molecular Biology
 DT Journal
 LA English
 AB Several glycine ribonucleotide analogs have been prepared and evaluated as substrates and/or inhibitors of glycine ribonucleotide transformylase from chicken liver. The side chain modified analogs, in which the glycine side chain, R = CH₂NH₂, has been replaced by R = CH₂NHCH₃ and R = CH₂CH₂NH₂, are substrates, with V/K (relative intensity) of 2.4% and 16.3%, resp. Several carbocyclic analogs of glycine ribonucleotide, including the phosphonate derivative of carbocyclic glycine ribonucleotide, did not serve as substrates, but were inhibitors of the enzyme, competitive against glycine ribonucleotide, with K_i values ranging from 7.4 to 23.6 times the K_m for glycine ribonucleotide. However, the O-phosphonate analog of carbocyclic glycine ribonucleotide did support enzymic activity, with V/K (relative intensity) of 0.8%. In addition, glycine ribonucleoside was neither a substrate for, nor an inhibitor of, glycine ribonucleotide transformylase. Furthermore, α-glycine ribonucleotide had no effect on enzyme activity. These studies have begun to define the structural features of the nucleotide substrate required to support enzymic activity.
 IT 127061-46-5
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (substrate and inhibitor specificity of glycine ribonucleotide transformylase from chicken liver)
 RN 127061-46-5 CAPLUS
 CN 2-Cyclopentene-1-carboxylic acid, 4-(acetamino)-, methyl ester, (1S-cis)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L43 ANSWER 6 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1995:579673 CAPLUS
 DN 123:286496
 TI Biocatalytical transformations. VI. The 4-acetamido-cyclopent-2-ene carboxylate route revisited: synthesis of (+)- and (-)- aristeromycin
 AU Csuk, Rene; Doerr, Petra
 CS Pharm.-Chem. Inst., Univ. Heidelberg, Heidelberg, D-69120, Germany
 SO Tetrahedron (1995), 51(20), 5789-98
 CODEN: TETRAB; ISSN: 0040-4020
 PB Elsevier
 DT Journal
 LA English
 AB Enantiomerically pure(+)- as well as (-)-aristeromycin can be synthesized starting from (+)- or (-)-Bu (or hexyl) 4-acetamido-cyclopent-2-ene carboxylate; these carboxylates are easily obtained from their corresponding racemates by hydrolysis with the lipase from *Candida rugosa*.
 IT 155836-24-1P
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)
 (enzymic hydrolysis of acetamidocyclopentene carboxylate in preparation of aristeromycin)
 RN 155836-24-1 CAPLUS
 CN 2-Cyclopentene-1-carboxylic acid, 4-(acetamino)-, butyl ester, (1S-cis)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 127061-46-5P
 RL: BPN (Biosynthetic preparation); RCT (Reactant); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)
 (enzymic hydrolysis of acetamidocyclopentene carboxylate in preparation of aristeromycin)
 RN 127061-46-5 CAPLUS
 CN 2-cyclopentene-1-carboxylic acid, 4-(acetylamino)-, methyl ester, (1S-cis)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L43 ANSWER 7 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1995:36909 CAPLUS
 DN 123:83906
 TI The enantioselective synthesis of an important intermediate to the antiviral, (-)-carbovir
 AU Handa, Sheetal; Earlam, George J.; Geary, Phillip J.; Hawes, John E.; Phillips, Gareth T.; Pryce, Robert J.; Ryback, George; Shears, Jeremy H.
 CS Department of Chemistry, King's College London, London, WC2R 2LS, UK
 SO Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry (1972-1999) (1994), (14), 1885-6
 CODEN: JCPRB4; ISSN: 0300-922X
 DT Journal
 LA English
 OS CASREACT 123:83906
 AB Two new routes to the important intermediate (-)-I (R = NHAc) for the carbocyclic-based nucleosides are reported. The intermediate (-)-I (R = NHAc) has also been synthesized in high enantiomeric excess via an enzymic resolution of the racemic amide (+)-I (R = NHAc) or an enzymic enantiotopic hydrolysis of the meso diester I (R = CO2Me).
 IT 127061-46-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (enantioselective synthesis of amide cyclopentadiene as carbovir synthon)
 RN 127061-46-5 CAPLUS
 CN 2-cyclopentene-1-carboxylic acid, 4-(acetylamino)-, methyl ester, (1S-cis)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L43 ANSWER 8 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1994:436093 CAPLUS
 DN 121:36093
 TI Biocatalytical transformations. IV. Enantioselective enzymic hydrolyses of building blocks for the synthesis of carbocyclic nucleosides
 AU Csuk, Rene; Doerr, Petra
 CS Pharm.-Chem. Inst., Univ. Heidelberg, Heidelberg, D-69120, Germany
 SO Tetrahedron: Asymmetry (1994), 5(2), 269-76
 CODEN: TASYE3; ISSN: 0957-4166
 DT Journal
 LA English
 OS CASREACT 121:36093
 AB Enantiomerically pure alkyl acetamidocyclopentenecarboxylates I (R = NHAc, R1 = R2 = H, R3 = CO2Me; R = R3 = H, R1 = NHAc, R2 = CO2Me) are obtained from their corresponding racemates by hydrolysis with PLE or the lipase from Candida cylindracea.
 IT 155836-24-1P 155836-25-2P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and sequential hydrolysis and esterification of)
 RN 155836-24-1 CAPLUS
 CN 2-cyclopentene-1-carboxylic acid, 4-(acetylamino)-, butyl ester, (1S-cis)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 155836-25-2 CAPLUS

CN 2-Cyclopentene-1-carboxylic acid, 4-(acetylamino)-, hexyl ester, (1S-cis)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 127061-46-5P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of, as building block for the synthesis of carbocyclic nucleosides)

RN 127061-46-5 CAPLUS

CN 2-Cyclopentene-1-carboxylic acid, 4-(acetylamino)-, methyl ester,
(1S-cis)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L43 ANSWER 9 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1992:236065 CAPLUS

DN 116:236065

TI Potential use of carbocyclic nucleosides for the treatment of AIDS:
chemo-enzymic syntheses of the enantiomers of carbovir

AU Evans, Chris T.; Roberts, Stanley M.; Shoberu, Karoline A.; Sutherland,
Alan G.

CS Enzymatix Ltd., Cambridge, SW4 4WE, UK

SO Journal of the Chemical Society, Perkin Transactions 1: Organic and
Bio-Organic Chemistry (1972-1999) (1992), (5), 589-92

CODEN: JCPRB4; ISSN: 0300-922X

DT Journal

LA English

AB Lactam azabicycloheptenone I, derived by whole cell enantioselective
hydrolysis of the racemate was converted into (-)-carbovir in 10 steps.
Lipase-catalyzed acetylation of 4-cis-hydroxycyclopent-2-enylmethyl
triphenylmethyl ether afforded the optically pure ester II (R = CPh3, R1
= Ac) (III) and the alc. II (R = H, R1 = CPh3). III was converted into
(+)-carbovir in 3 steps.


IT 127061-46-5P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(preparation and reduction of)

RN 127061-46-5 CAPLUS

CN 2-Cyclopentene-1-carboxylic acid, 4-(acetylamino)-, methyl ester,
(1S-cis)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L43 ANSWER 10 OF 10 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1990:217364 CAPLUS

DN 112:217364

TI Synthesis of the two enantiomers of the carbocyclic analog of nicotinamide
ribose and analysis of their biological properties

AU Ikbal, Mohamed; Cerceau, Claude; Le Goffic, Francois; Sicsic, Sames

CS CERCOA, CNRS, Thiais, 94320, Fr.

SO European Journal of Medicinal Chemistry (1989), 24(4), 415-20

CODEN: EJMCA5; ISSN: 0223-5234

DT Journal

LA French

OS CASREACT 112:217364


AB Enantiomers of the carbocyclic analog of nicotinamide ribose I were prepared
via an enzymic resolution of the precursor (\pm)-II using pig liver
esterase. (-)-I possessed good and highly specific bactericidal and
fungicidal activities. In vivo competition expts. between (-)-I and
intermediate mols. of the pyridine nucleotide cycle along with its
inhibitory behavior against 2 key enzymes of the NAD⁺ metabolism were
performed and suggested that the target of (-)-I could be one of the
enzymes involved in NAD⁺ metabolism

IT 127061-46-5P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and hydroxylation by osmium tetroxide)

RN 127061-46-5 CAPLUS
CN 2-Cyclopentene-1-carboxylic acid, 4-(acetylamino)-, methyl ester,
(1S-cis)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> FIL STNGUIDE
FILE 'STNGUIDE' ENTERED AT 21:43:39 ON 10 JUN 2006
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Jun 9, 2006 (20060609/UP).

=> d his